

## TECHNICAL NOTE

# Influence of *Houttuynia Cordata* Powder on The Growth Performance of Ducks and The Impact of $\text{AlCl}_3$ Treatment on Ammonia flux in Duck Litter

Tae-Ho Chung, In-Hag Choi\*

Department of Companion Animal & Animal Resources Science, Joongbu University, Geumsan-gun 312-702, South Korea

## Abstract

The effects of *Houttuynia cordata* powder on the growth performance of ducks were investigated. Ninety ducks were assigned into one of three dietary treatments as a completely randomized design for 6 weeks: feeds supplemented with 1% or 2% *H. cordata* and a control group. No significant difference was observed in feed conversion among the three groups ( $p > 0.05$ ), but addition of *H. cordata* had a significantly positive effect ( $p < 0.05$ ) on initial and final body weight, weight gain, and feed intake of the ducks. Furthermore, the effects of chemical treatment (comprising 50 g and 100 g aluminum chloride [ $\text{AlCl}_3$ ] per kilogram litter) on the ammonia ( $\text{NH}_3$ ) flux in duck litters were also investigated. Duck litter was treated with  $\text{AlCl}_3$  at a depth of 8 cm by top-dressing; this resulted in a significant difference on  $\text{NH}_3$  flux ( $p < 0.05$ ) during the experimental period (but not at 2 weeks).  $\text{NH}_3$  flux at 6 weeks were reduced by 25.4% and 37.5% by treatment with 1% and 2% *H. cordata*, respectively, compared with the control groups. In conclusion, enriching the diets of the ducks with 2% *H. cordata* and adding 100 g  $\text{AlCl}_3$  to their litter has beneficial effects on increasing their growth performance and reducing  $\text{NH}_3$  flux in their environment.

**Key words** : *Houttuynia cordata* powder, Aluminum chloride, Growth performance, Ammonia, Duck litter

## 1. Introduction

Antibiotic growth promoters (AGPs) are used in animal diets to enhance livestock production and economic viability, as well as to prevent subclinical diseases. However, the use of antibiotics in animal feed has been prohibited in many countries due to the rise of pathogen resistance to antibiotics. In January 2006, the European Union banned the use of AGPs in animal feed except for three antibiotics (salinomycin-Na, flavophospholipol, avilamycin) (Wenk, 2003; Catalá-Gregori et al., 2008). Consequently, many

alternative approaches that have the potential to maintain animal health and productivity in the livestock industry have been investigated. Among these, medicinal herbs are considered a potential candidate for antibiotic alternatives.

*Houttuynia cordata* (*H. cordata*) is traditionally used as an aromatic medicinal herb because of its antioxidant, antimicrobial, antiviral, and anti-inflammatory activities. It is distributed widely in China, Korea, India, and Taiwan (Hayashi et al., 1995; Chen et al., 2003; Lu et al., 2006). *H. cordata* is known to have beneficial effects on livestock productivity,

Received 20 September, 2015; Revised 13 October, 2015;

Accepted 15 October, 2015

\*Corresponding author : In-Hag Choi, Department of Companion Animal & Animal Resources Science, Joongbu University, Geumsan-gun 312-702, South Korea  
Phone: +82-41-750-6284  
E-mail: wiew@chol.com

© The Korean Environmental Sciences Society. All rights reserved.  
© This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

nutrient digestibility, and mast cell-mediated inflammation (Kim et al, 2007; Yan et al., 2011). In addition, the major components in *H. cordata* are essential oils, alkaloids, and flavonoids that possess antioxidant properties and free radical scavenging capacities (Fu et al., 2013).

Another challenge faced by the livestock industry is the regulation of ammonia ( $\text{NH}_3$ ) levels in poultry litter. The buildup of  $\text{NH}_3$  in the environment has inevitably provoked various concerns about environmental, human, and poultry health. However, litter is a valuable resource for improving crop production. Thus, recognizing and understanding its significance is central to good litter management in poultry houses. Management practices could be refined to mitigate potential negative environmental impacts of litter on air and soil. For example, chemical amendments to animal manure or litter have been shown to reduce  $\text{NH}_3$  levels effectively; a study by Smith et al. (2001) demonstrated that swine manure treated with  $\text{AlCl}_3$  emitted less  $\text{NH}_3$ .

Therefore, the objective of the present study was to evaluate the beneficial effects of *H. cordata* on the growth performance in ducks and furthermore examine how ammonia emissions are affected by addition of  $\text{AlCl}_3$  to duck litter?

## 2. Material and methods

### 2.1. Leaf preparation

*H. cordata* leaves and stems were obtained from a herbal medicine market (Daegu, South Korea). They were initially air-dried for 12 h at room temperature and then oven-dried at 50°C for two consecutive days and subsequently ground to a fine texture. The resulting powder was stored in airtight plastic bags until further processing.

### 2.2. Experimental design and birds

All experimental protocols were performed in

compliance with the animal care guidelines of animal policy at Gilhong farm (Geochang, South Korea). Ninety ducks (one-day-old pekin, 45 male and 45 female) were assigned to one of three dietary treatments in a completely randomized design for a 6-week experimental period: two treatment groups (T1 [1% *H. cordata*] and T2 [2% *H. cordata*]) and one control group (Control). After a brooding period of 1 week, each treatment group was subdivided into three replicates of 10 ducks (5 male and 5 female) per pen. Starter diets were provided from day 1 to 21 and comprised 21% crude protein, 2.5% crude fat, 8% crude fiber, 9% crude ash, 0.40% Ca, and 1.50% P. Finisher diets provided from day 22 to 42 consisted of 17% crude protein, 2.5% crude fat, 8% crude fiber, 9% crude ash, 0.40% Ca, and 1.0% P. A feeder and a drinker were placed in each pen and the ducks had access to feed and water filled *ad lib*. Approximately 8 cm of litter comprising rice hulls and duck manure was deposited over concrete flooring. Ventilation and temperature in the duck houses were automatically regulated. Ducks were weighed at 8 and 42 days of age, and the average weights were recorded to determine growth performance. Feed intake was also recorded at each feed change interval during the experimental period. Body weight gain and feed intake were used to calculate feed conversion ratio (FCR).

### 2.3. Chemical treatment and ammonia measurement

The duck litter with rice hulls was chemically treated by top-dressing. The three treatments consisted of control, T1 (50 g  $\text{AlCl}_3/\text{kg}$  litter) and T2 (100 g  $\text{AlCl}_3/\text{kg}$  litter). Aluminum chloride ( $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ ) was purchased from Samchun Chemicals Company (Pyeongtaek, South Korea). Ammonia emissions from duck litter with rice hulls were measured weekly (2 weeks through 6 weeks) at 4 random sites in each pen by using a multi-gas analyzer (Yes Plus LGA, Critical Environment Technologies Canada