

decreased in catagen hair follicles, especially free fatty acids and was ester. These results suggest that changes of IHL might be related to hair cycle.

Key Words: Hair cycle, Integral hair lipid

## P170

### Epidemiologic characteristics among juvenile alopecia areata patients : A retrospective study of 157 patients from Korea

Department of Dermatology and Institute of Hair and Cosmetic Medicine, Yonsei University Wonju College of Medicine, Wonju, Korea.

**Na Young Yoon, Bo Kyung Kim, Sung-yul Lee, Seok-Yong Ahn, Won-Soo Lee**

Alopecia areata (AA) is a common cause of nonscarring alopecia and considered an autoimmune disease with undetermined pathogenesis. To study the clinical and epidemiologic profile of juvenile alopecia, we performed a survey in which a total of 157 patients younger than 18 years old were enrolled. The male: female ratio was 1:1.2. The median age of onset was 10.6 years old and the mean disease duration was 5.9 months. 28%(44 cases) patients had past medical histories(atopic diseases and autoimmune diseases) and the most common disease was atopic dermatitis. The early onset group showed more severe types. In this survey, 79.9% (125 cases) patients had common type showing one to four coin shaped alopecia patches. 20.1% (32 cases) patients had severe type showing large and severe patchy alopecia (9%, 18cases), alopecia totalis (1%, 2cases) and alopecia universalis (10%, 12cases). In this study, we checked several lab findings including Hb, Hct, BUN/Cr, AST/ALT, fT4, and ANA. There were no noticeable differences between the various lab tests. This retrospective study reveals some clinical characteristics of Korean juvenile alopecia areata patients.

Key Words: Juvenile alopecia areata, Korea

## P171

### Epidemiologic characteristics and comorbidity profiles among alopecia areata patients - a retrospective study of 871 Korean patients

Department of Dermatology and Institute of Hair and Cosmetic Medicine, Yonsei University Wonju College of Medicine, Wonju, Korea.

**Noo Ri Lee, Sung-yul Lee, Seok-Yong Ahn, Won-Soo Lee**

Alopecia areata (AA) is an autoimmune disease that presents as patchy, nonscarring hair loss, affecting about 2% of the population. AA is thought to occur in association with autoimmune diseases such as thyroid disorders, atopic dermatitis. Only a few studies have been investigated about clinical profiles of Asian AA patients. In this study, we have performed retrospective study of 871 patients who were diagnosed as alopecia areata in the last 10 years at Yonsei Wonju Christian Hospital, Department of Dermatology. Male to female ratio was 1:1.01, the mean age at presentation was 34.1 years. The peak age was in the thirties in male, and forties in female. The frequencies of the following associated diseases were: hypertension, 3.4%; diabetes mellitus, 2.2%; hypercholesterolemia, 0.9%; thyroid disease, 4.4%; atopic dermatitis, 5.2%. These findings are similar to those reported in a study of 219 Singapore patients. To evaluate other possible associated abnormalities, we have done serologic tests on the first day of visit to the clinic, including anemia test, thyroid function test, glucose level, lipid profiles, liver enzymes, male hormone levels and autoimmune test. As results, no remarkable abnormalities revealed in the laboratory results. Although there are some limits of this study because of the retrospective design, this study has an importance that epidemiologic research is done in a large group of Asian AA patients.

Key Words: Alopecia areata, Clinical profile, Asian

## P172

### The effect on the hair growth of combination therapy of microneedle roller and hair tonic with and without phytosphingosine-1-phosphate in female pattern hair loss

Department of Dermatology, Asan Medical Center, University of Ulsan College of Medicine, Seoul

**Sang Min Lee, Gyeong-hun Park, Chong Hyun Won, Sung Eun Chang, Mi Woo Lee, Jee Ho Choi, Kee Chan Moon**

Phytosphingosine-1-phosphate is an analogue of sphingosine-1-phosphate that is a kind of lipid mediator known to regulate angiogenesis, cell migration and proliferation. A

16-week double-blind randomized controlled clinical trial was conducted to evaluate the effect of the combination therapy of microneedle roller and hair tonic with and without phytosphingosine-1-phosphate on the female pattern hair loss in 36 women. We evaluated the change of total, terminal (more than 40  $\mu\text{m}$  in diameter) and vellus (less than 40  $\mu\text{m}$ ) hair counts with phototrichogram at baseline and 8 and 16 weeks after treatment start. Mean change of hair counts from baseline to 8 and 16 weeks was an increase of 4.2 and 8.8 / $\text{cm}^2$  respectively in phytosphingosine-1-phosphate containing hair tonic group, and 4.1 and 5.2 / $\text{cm}^2$  in control hair tonic group. In addition, mean increases of terminal hair counts at 8 and 16 weeks were 7.1 and 6.8 / $\text{cm}^2$  respectively in phytosphingosine-1-phosphate containing group, and 0.3 and 2.7 / $\text{cm}^2$  in control group. These results suggest that phytosphingosine-1-phosphate may further stimulate the growth of hairs, especially terminal hairs, compared with microneedle roller therapy.

Key Words: Phytosphingosine-1-phosphate, Microneedle, Female pattern hair loss, Androgenetic alopecia

## P173

### The effect of red ginseng on the growth of cultured human hair follicles

Department of Dermatology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea, <sup>1</sup>Asan Institute for Life Sciences, Seoul, Korea

**Joon-min Jung, Gyeong-hun Park, Chong-Hyun Won, Ki-young Park<sup>1</sup>, Sang-Min Lee, Sung-Eun Chang, Mi-Woo Lee, Jee-Ho Choi, Kee-Chan Moon, Hong-il Cho<sup>1</sup>**

Several recent murine studies showed that ginseng and its bioactive components, saponins, may be beneficial to hair growth promotion. However, the effects on human hair follicles have not yet been elucidated sufficiently. To investigate the effect of red ginseng and its saponins on the hair growth. We determined the proliferation of human dermal papilla cells (hDPCs) using MTT assay. The effect on cultured human hair follicles was assessed by the proliferation of hair matrix keratinocytes with immunofluorescence staining of Ki67. In addition, we investigated the signal changes of hDPCs and the effect on hair growth in C57BL/6 mice. The red ginseng extract, ginsenoside Rb1 and Rg1 increased the proliferation of hDPCs in MTT assays, and upregulated the proliferation of hair matrix keratinocytes in

the cultured human hair follicles. Both red ginseng extract and ginsenoside Rb1 increased the phosphorylation of ERK1/2 in hDPCs. Furthermore, the subcutaneous injection of 3% red ginseng extract resulted in the hair growth similar to that by topical application of 2% minoxidil solution. The red ginseng extract and its ginsenosides may enhance the proliferation of hDPCs, activate the ERK signaling pathway in hDPCs, and upregulate the proliferation of hair matrix keratinocytes. These results suggest that red ginseng may be beneficial for human hair growth promotion.

Key Words: Red ginseng

## P174

### Efficacy of home treatment using diphencycloproprone for alopecia areata : Focused on patients' safety, convenience and economic feasibility

Department of Dermatology, Chonbuk National University Medical School

**Ki-Hun Song, Kyung Hwa Nam, Jin Park, Seok Kweon Yun, Han Uk Kim**

Immunotherapy using diphencycloproprone (DPCP) is an effective treatment modality for alopecia areata, although there are some troubling side effects as DPCP is a potent contact sensitizer. Hence, most of DPCP immunotherapy is directly conducted by dermatologists at a hospital weekly. However, since alopecia areata usually requires long-term treatment, frequent visiting to a hospital could be a considerable loss of time and money for patients, which might lead to decline in quality of life. To mitigate inconveniences, we performed home treatment which is applying DPCP solution by patients themselves or guardians at home. We retrospectively studied 87 patients with alopecia areata who received home treatment. Sensitization was done by dermatologists at the hospital before starting home treatment. They received sufficient educations regarding application methods and possible side effects. All patients were given the optimal concentration of DPCP solution at each visit. Follow-up at the hospital was done monthly, and the authors assessed clinical features such as regrowth of hair using SALT score, side effects of DPCP immunotherapy and patient's quality of life. The results of this study were compared with that of previously published papers in which DPCP immunotherapy was done at hospitals.

Key Words: Alopecia areata, Diphencycloproprone, Home